

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12Q 1/68, C12N 15/00, 15/09, 15/63, 15/86		A1	(11) International Publication Number: WO 00/55375 (43) International Publication Date: 21 September 2000 (21.09.00)																		
(21) International Application Number: PCT/US00/07285 (22) International Filing Date: 17 March 2000 (17.03.00) (30) Priority Data: <table><tr><td>60/124,916</td><td>17 March 1999 (17.03.99)</td><td>US</td></tr><tr><td>60/124,808</td><td>17 March 1999 (17.03.99)</td><td>US</td></tr><tr><td>60/149,639</td><td>17 August 1999 (17.08.99)</td><td>US</td></tr><tr><td>60/157,247</td><td>1 October 1999 (01.10.99)</td><td>US</td></tr><tr><td>60/167,824</td><td>29 November 1999 (29.11.99)</td><td>US</td></tr><tr><td>60/182,711</td><td>15 February 2000 (15.02.00)</td><td>US</td></tr></table> (71) Applicant: ALPHAGENE, INC. [US/US]; 260 West Cummings Park, Woburn, MA 01801 (US). (72) Inventors: VALENZUELA, Dario; 1081 Hill Road, Boxborough, MA 01719-1010 (US). YUAN, Olive; 292 Mystic Street, Arlington, MA 02174 (US). HOFFMAN, Heidi; 90 Houghton Mill Road, Lunenburg, MA 01462 (US). HALL, Jeff; 4 Alderwood Drive, Stratham, NH 03885 (US). RAPIEJKO, Peter; 63 Old Grafton Road, Upton, MA 01568 (US). (74) Agent: SPRUNGER, Suzanne, A.; American Home Products Corporation, Patent & Trademark Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US).			60/124,916	17 March 1999 (17.03.99)	US	60/124,808	17 March 1999 (17.03.99)	US	60/149,639	17 August 1999 (17.08.99)	US	60/157,247	1 October 1999 (01.10.99)	US	60/167,824	29 November 1999 (29.11.99)	US	60/182,711	15 February 2000 (15.02.00)	US	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
60/124,916	17 March 1999 (17.03.99)	US																			
60/124,808	17 March 1999 (17.03.99)	US																			
60/149,639	17 August 1999 (17.08.99)	US																			
60/157,247	1 October 1999 (01.10.99)	US																			
60/167,824	29 November 1999 (29.11.99)	US																			
60/182,711	15 February 2000 (15.02.00)	US																			
(54) Title: SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM																					
(57) Abstract Novel polynucleotides and the proteins encoded thereby are disclosed.																					

#17
09/651,563 A

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Asn Lys His Ser Arg Ala Cys Gln Gln Phe Leu Lys Gln Cys Gln Leu
 100 105 110

Arg Ser Phe Ala Leu Pro Leu
 115


<210> 85
 <211> 1176
 <212> DNA
 <213> Homo sapiens

<400> 85
 aatctctttt taaataagta ttttgttgag gttgaagaat tgctggcaat taaaagaata 60
 gagctaatta tggctttcat cattcattca tgtatttatt gagcacctac ttattatggt 120
 gctcaacact tgttactgca agctacetta atttcccaag agtgggtgct tactctgttt 180
 tttctgatat ggtcttccaa tcagtgtgtg taacatacct gttgtttatc agccattgta 240
 ggtggctgtg tctgttgcac catcataaga agtttaagct ttgtgctctg ataaattgtg 300
 ttctgttaaa gaggttagta ggatgaaaac agcaaaacaa taattttttc aacaaattgt 360
 aaattataag aaaaagagtt ggtttgtgta caacaatttt aatgattccc ttgttcattt 420
 ttgctgtgaa atgcactgaa aaaaatcttc aaaatgagtt atagtccctg tgttgggaaa 480
 attgacaaat aataaaacta gagaacaaac aataatgctt ctgtttcttt tacgaatgga 540
 gagagaaaagt ttatattcag tagagttatt gccctgttca tttgagaggg gcatggattt 600
 tctgtttaag tcttccagga atcttcagct aggtggtaaa ttttaataaga gtttctaaaa 660
 attgaaatgt ttaactttta aatattctgg agatagaaga agaataataa atgaaaccag 720
 gctgatctgc atgcagtggc atttacaact aactgatcac aaccaattat agattcctta 780
 ttttgttat tgtgaggcag agtctgactc tgtcaccag gatggagtac agtgcatagc 840
 tcaactcagt cttagacctc caggctaaac caatcattcc acttcacctt cccaagtagc 900
 tgagaccaca ggcacacaac accacaacca gctgattgtt gtactgttgc tcaagcagtc ctcccacctc 1020
 agcctcccaa agtactggga ttacaggcgt gaggtacctc gcttcttctt ccttcttctt 1080
 ttttcttctt cactgtctta acttgattag ccttcttctt agaaataaaa 1140
 aaaaattttt aaaaaaaaaa aaaaaaaaaa aaaaaa 1176

<210> 86
 <211> 78
 <212> PRT
 <213> Homo sapiens

<400> 86
 Met Tyr Leu Leu Ser Thr Tyr Leu Leu Trp Cys Ser Thr Leu Val Thr
 1 5 10 15
 Ala Ser Tyr Leu Asn Phe Pro Arg Val Val Pro Tyr Ser Val Phe Ser
 20 25 30
 Asp Met Val Phe Gln Ser Val Cys Val Thr Tyr Leu Leu Phe Ile Ser
 35 40 45
 His Cys Arg Trp Leu Cys Leu Leu His His His Lys Lys Phe Lys Leu
 50 55 60
 Cys Ala Leu Ile Asn Cys Val Leu Leu Lys Arg Leu Val Gly
 65 70 75

<210> 87
 <211> 1476
 <212> DNA
 <213> Homo sapiens



<400> 87

gaggtctgag aggtgccatg gctgagtcac acctgctgca gtggctgctg ctgctgctgc 60
 ccacgtcttg tggcccaggc actgctgcct ggaccacctc atccttggcc tgtgcccagg 120
 gccctgagtt ctgggtgcaa agcctggagc aagcattgca gtgcagagcc ctaggggcatt 180
 gcctacagga agtctgggga catgtgggag ccgatgacct atgccaagag tgtgaggaca 240
 tcgtccacat ccttaacaag atggccaagg aggccatttt ccaggatctc tccgagcagc 300
 aattccccat tcctctcccc tattgctggc ctctgcaggg ctctgatcaa gcggatecaa 360
 gccatgattc ccaaggggtg gctagctgtg gcagtggccc aggtgtgccc cgtgggtacct 420
 ctggtggcgg gcggcatctg ccagtgcctg gctgagcgct actccgtcat cctgctcgac 480
 acgtctgtgg gccgcatgct gccccagctg gtctgccgcc tcgtccctccg gtgctccatg 540
 gatgacagcg ctggcccagg agaattggctg ccgcgagact ctgagtggca cctctgcatg 600
 tccgtgacca cccaggccgg gaacagcagc gagcaggcca taccacaggc aatgctccag 660
 gcctgtgttg gctcctggct ggacaggga aagtgcaggc aatttgtgga gcagcacacg 720
 cccagctgct tgaccctggg gccccagggg tgggatgccc acaccacctg ccaggccctc 780
 ggggtgtgtg ggaccatgtc cagccctctc cagtgtatcc acagccccga cctttgatga 840
 gaactcagct gtccagaaaa agacaccgtc ctttaaagtg ctgcagtatg gccagacgtg 900
 gtggctcaca cctgcaatcc cagcacctta ggaggccgag gcaggaggat ccttgaggtc 960
 aggagtctga gaccagcttc gccaacatgg tgaaccccca tttctactaa aaatacaaaa 1020
 aattagccaa gtgtgtgtgg atatgcctgt aatcccaact actcagaagg ccgaggcagg 1080
 agaattactt gaacgcagga gaactactgc agcccaggag gcagagggtg cagtgaagccg 1140
 agattgcacc actgcactcc agcctgggtg acagagcaag actccatctc agtaataaaa 1200
 taaataaata aaaagcgtg cagttagctg ggcctcacc tgaagtcagc gggcccaggc 1260
 ctacctact ctctcccttg gcagagaagc agacgtccat agtcctctc cctcacaagc 1320
 gctcccagcc tgccctccag ctgctgctct cccctcccag tctctactca ctgggatgag 1380
 gttaggtcat gaggacacca aaaaacctaa aataaacaag aagccaaaca agccttagct 1440
 tttcttaag acaaaaaaaaa aaaaaaaaa aaaaaa 1476

<210> 88


<211> 145

<212> PRT

<213> Homo sapiens

<400> 88

Met Ala Glu Ser His Leu Leu Gln Trp Leu Leu Leu Leu Leu Pro Thr
 1 5 10 15
 Leu Cys Gly Pro Gly Thr Ala Ala Trp Thr Thr Ser Ser Leu Ala Cys
 20 25 30
 Ala Gln Gly Pro Glu Phe Trp Cys Gln Ser Leu Glu Gln Ala Leu Gln
 35 40 45
 Cys Arg Ala Leu Gly His Cys Leu Gln Glu Val Trp Gly His Val Gly
 50 55 60
 Ala Asp Asp Leu Cys Gln Glu Cys Glu Asp Ile Val His Ile Leu Asn
 65 70 75 80
 Lys Met Ala Lys Glu Ala Ile Phe Gln Asp Leu Ser Glu Gln Gln Phe
 85 90 95
 Pro Ile Pro Leu Pro Tyr Cys Trp Pro Leu Gln Gly Ser Asp Gln Ala
 100 105 110
 Asp Pro Ser His Asp Ser Gln Gly Cys Ala Ser Cys Gly Ser Gly Pro
 115 120 125
 Gly Val Pro Arg Gly Thr Ser Gly Gly Gly Arg His Leu Pro Val Pro
 130 135 140




Gly
145

<210> 89
<211> 2243
<212> DNA
<213> Homo sapiens

<400> 89
agtagctttgt aatcaagtga aaatataact ttattttttta actctattac attttatttt 60
gtcatgtact aaaattattt ctgtattgct ttatataaaa acagtggcat ttagcactgg 120
cattgagact atagcacatc atttttgcca ttttcagtgc ttatattgtt aggttagaggc 180
tgccacttta ttagaatgca agccacaaaa atatcaattt tgtttttttt gttagggtgg 240
gtcttctttt ttctttccc tctctctttt ttaacaaat gccttcttat agaaaaactt 300
tctaagaggg aacaatttag aatggatatt ttgacgaatc ggcattgagtg taacagtgat 360
aacctgatct gtttgtttta aagattatta ccaagtgaat aattcagaat gaatagaatt 420
tacactaaca tgetatataa aatgttaag tctgatgctg tgaagcaat ctagtgttat 480
atttctacct cctcatttgt cttaattatt tggtaagtgg gattatgatg agtaactgga 540
ggggcttaga aacaaaaact ggtgaaaga gtatgcatga agaaaagctt ctttgataaa 600
tgtggagttc ttcattataa atatatattc atgaattcac agataagtac ttaagaaca 660
gacagtttac ttggcctana aatattttga tgtttactca aaaagtacct cttcagggtc 720
tgagaacatg gaaaagaatt gaggctttt aaatactttt tagaaagtaa tcataaagt 780
aaattgaatt tcaaacctat ttggcttctg ttttgtaac ctttgaacta tatgtatgtg 840
tataagggtg tacacatata tatatggcat ataacaagtg tacacatata cacataacaa 900
gtgtagaagt atatatataa tacatacact cactctgtct ggtataggtc aattttgaag 960
aactcccata agtttctgct gcttctccca taactgctgc caccaccatc agaattcata 1020
atcaaaccta acctttttgt ttggggcacc aaatctgaag acaaaattaa ttgaccagg 1080
taaacttcaa gctgctttct ttcttgaana cttaacgttt aacgtataat gtctgtttgg 1140
atactgttcc aaattgttga ttgcatgtgg ttaattgtgc attagagcac ttgcaattg 1200
cataattcat taatgttttg tgagcttgca tttgtgattt attggatgat cagactgaat 1260
tttgtaagat atcacattgt acatcttgcc tagatgtcga tgactgcaag taataataa 1320
gtttataatg aaactatcta caattcttgt tttagcacat ctgttatccg taaaacacct 1380
gtaactagct tttttaattt attatttgaa ttttaggata gcgaatcact aatttttagt 1440
tgctgaggtt ggcatttttag tgattattaa gcacttctgt cagtctttga aaaaagaacg 1500
tattttttgt gctttgaaga tctctgaaga atttctttta taatagaatg ggcattgatt 1560
gtaacagttt tatgtcaaat gatctgtgct gtagaaaaac attaacctt gttcaaaaaa 1620
gaaatggata aacttggcct ttctaagtgg taagaatgac ctgtcactat aatatactgt 1680
atgtttacat tttattttaa ttttatctct tatgtatagg gtgataacct tcccagaaa 1740
caacagtgat tgcgattgtt ttctagaac ttctttaag tgccacattt ggcagtacaa 1800
atgagcttga gtgtaatagc ccagagattt atatatagtt gaatgtctaa aatggtaaaa 1860
tgtgccactg tgcgaagtta cagtggctta tgtttttcat agtaattcaa atgaacttcc 1920
tatttttgat agtaaatgtc atttaatagt atacttgcga tttgagctc actgcaaaat 1980
tagtgcagag gagaaaaaaa tttttaatgt aatcttgatt ttacctcata tactgtacat 2040
tccaaaaact cttaactttt taaagattat agatacacta ccaacatat caccttaaaa 2100
ttgtataagg ctgaatgaac ttcatacaaa tgaaaaaaat ctcataaaaa tacataaact 2160
atgtagcaaa agtatctgta aaatccatgg aaataaaaag ttgtatcatt ctttttgaaa 2220
aaaaaaaaaa aaaaaaaaaa aaa 2243

<210> 90
<211> 61
<212> PRT
<213> Homo sapiens

<400> 90
Met Gln Ala Thr Lys Ile Ser Ile Leu Phe Phe Leu Leu Gly Trp Val
1 5 10 15
Phe Phe Phe Ser Phe Pro Leu Ser Phe Phe Asn Lys Cys Leu Leu Ile



What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
 - (a) the nucleotide sequence of SEQ ID NO:1;
 - (b) the nucleotide sequence of SEQ ID NO:1 from nucleotide 27 to nucleotide 260;
 - (c) the nucleotide sequence of SEQ ID NO:1 from nucleotide 72 to nucleotide 260;
 - (d) the nucleotide sequence of the full-length protein coding sequence of clone vc62_1 deposited with the ATCC under accession number 207114;
 - (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc62_1 deposited with the ATCC under accession number 207114;
 - (f) the nucleotide sequence of a mature protein coding sequence of clone vc62_1 deposited with the ATCC under accession number 207114;
 - (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc62_1 deposited with the ATCC under accession number 207114;
 - (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
 - (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
 - (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
 - (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:1.

2. The polynucleotide of claim 1 wherein said polynucleotide is operably linked to at least one expression control sequence.
3. A host cell transformed with the polynucleotide of claim 2.
4. The host cell of claim 3, wherein said cell is a mammalian cell.
5. A process for producing a protein encoded by the polynucleotide of claim 2, which process comprises:
 - (a) growing a culture of a host cell in a suitable culture medium, wherein the host cell has been transformed with the polynucleotide of claim 2; and
 - (b) purifying said protein from the culture.
6. A protein produced according to the process of claim 5.
7. An isolated polynucleotide encoding the protein of claim 6.
8. The polynucleotide of claim 7, wherein the polynucleotide comprises the cDNA insert of clone vc62_1 deposited with the ATCC under accession number 207114.
9. A protein comprising an amino acid sequence selected from the group consisting of:
 - (a) the amino acid sequence of SEQ ID NO:2;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vc62_1 deposited with the ATCC under accession number 207114;the protein being substantially free from other mammalian proteins.
10. The protein of claim 9, wherein said protein comprises the amino acid sequence of SEQ ID NO:2.

11. A composition comprising the protein of claim 9 and a pharmaceutically acceptable carrier.

12. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:3;
- (b) the nucleotide sequence of SEQ ID NO:3 from nucleotide 6 to nucleotide 1325;
- (c) the nucleotide sequence of SEQ ID NO:3 from nucleotide 99 to nucleotide 1325;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp10_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp10_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp10_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp10_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:3.

13. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:4;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vp10_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

14. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:5;
- (b) the nucleotide sequence of SEQ ID NO:5 from nucleotide 149 to nucleotide 322;
- (c) the nucleotide sequence of SEQ ID NO:5 from nucleotide 200 to nucleotide 322;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp11_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp11_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp11_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp11_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:5.

15. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:6;

(b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp11_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

16. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:7;

(b) the nucleotide sequence of SEQ ID NO:7 from nucleotide 288 to nucleotide 629;

(c) the nucleotide sequence of SEQ ID NO:7 from nucleotide 363 to nucleotide 629;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vp13_1 deposited with the ATCC under accession number 207114;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp13_1 deposited with the ATCC under accession number 207114;

(f) the nucleotide sequence of a mature protein coding sequence of clone vp13_1 deposited with the ATCC under accession number 207114;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp13_1 deposited with the ATCC under accession number 207114;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:8;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:7.

17. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:8;

(b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp13_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

18. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:9;

(b) the nucleotide sequence of SEQ ID NO:9 from nucleotide 11 to nucleotide 298;

(c) the nucleotide sequence of SEQ ID NO:9 from nucleotide 149 to nucleotide 298;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp16_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp16_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp16_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp16_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:9.

19. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vp16_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

20. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:11;
- (b) the nucleotide sequence of SEQ ID NO:11 from nucleotide 257 to nucleotide 607;
- (c) the nucleotide sequence of SEQ ID NO:11 from nucleotide 479 to nucleotide 607;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp21_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp21_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp21_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp21_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:12;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:11.

21. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vp21_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

22. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:13;
- (b) the nucleotide sequence of SEQ ID NO:13 from nucleotide 163 to nucleotide 477;
- (c) the nucleotide sequence of SEQ ID NO:13 from nucleotide 238 to nucleotide 477;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp22_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp22_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp22_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp22_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:14;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:13.

23. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:14;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vp22_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

24. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:15;
- (b) the nucleotide sequence of SEQ ID NO:15 from nucleotide 58 to nucleotide 624;
- (c) the nucleotide sequence of SEQ ID NO:15 from nucleotide 106 to nucleotide 624;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq2_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq2_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq2_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq2_1 deposited with the ATCC under accession number 207114;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:16;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:15.

25. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:16;

(b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq2_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

26. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:17;

(b) the nucleotide sequence of SEQ ID NO:17 from nucleotide 773 to nucleotide 1090;

(c) the nucleotide sequence of SEQ ID NO:17 from nucleotide 842 to nucleotide 1090;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq3_1 deposited with the ATCC under accession number 207114;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq3_1 deposited with the ATCC under accession number 207114;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq3_1 deposited with the ATCC under accession number 207114;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq3_1 deposited with the ATCC under accession number 207114;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:18;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:17.

27. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:18;

(b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq3_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

28. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:19;
- (b) the nucleotide sequence of SEQ ID NO:19 from nucleotide 96 to nucleotide 275;
- (c) the nucleotide sequence of SEQ ID NO:19 from nucleotide 159 to nucleotide 275;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq5_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq5_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq5_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq5_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:19.

29. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:20;

- (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vq5_1 deposited with the ATCC under accession number 207114; the protein being substantially free from other mammalian proteins.

30. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:21;
- (b) the nucleotide sequence of SEQ ID NO:21 from nucleotide 176 to nucleotide 340;
- (c) the nucleotide sequence of SEQ ID NO:21 from nucleotide 230 to nucleotide 340;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq6_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq6_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq6_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq6_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:21.

31. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vq6_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

32. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:23;
- (b) the nucleotide sequence of SEQ ID NO:23 from nucleotide 29 to nucleotide 1111;
- (c) the nucleotide sequence of SEQ ID NO:23 from nucleotide 167 to nucleotide 1111;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vr1_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vr1_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vr1_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vr1_1 deposited with the ATCC under accession number 207114;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:24;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:23.

33. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:24;

(b) a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24; and

(c) the amino acid sequence encoded by the cDNA insert of clone vr1_1 deposited with the ATCC under accession number 207114; the protein being substantially free from other mammalian proteins.

34. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:25;

(b) the nucleotide sequence of SEQ ID NO:25 from nucleotide 13 to nucleotide 513;

(c) the nucleotide sequence of the full-length protein coding sequence of clone vc63_1 deposited with the ATCC under accession number 207115;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc63_1 deposited with the ATCC under accession number 207115;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:26;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:25.

35. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:26;

(b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc63_1 deposited with the ATCC under accession number 207115;

the protein being substantially free from other mammalian proteins.

36. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:27;

(b) the nucleotide sequence of SEQ ID NO:27 from nucleotide 79 to nucleotide 345;

(c) the nucleotide sequence of SEQ ID NO:27 from nucleotide 130 to nucleotide 345;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vb25_1 deposited with the ATCC under accession number PTA-362;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb25_1 deposited with the ATCC under accession number PTA-362;

(f) the nucleotide sequence of a mature protein coding sequence of clone vb25_1 deposited with the ATCC under accession number PTA-362;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb25_1 deposited with the ATCC under accession number PTA-362;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:28;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:27.

37. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:28;

(b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb25_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

38. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:29;
- (b) the nucleotide sequence of SEQ ID NO:29 from nucleotide 72 to nucleotide 236;
- (c) the nucleotide sequence of SEQ ID NO:29 from nucleotide 150 to nucleotide 236;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb27_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb27_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb27_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb27_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:30;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:29.

39. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;

- (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vb27_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

40. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:31;
- (b) the nucleotide sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 884;
- (c) the nucleotide sequence of SEQ ID NO:31 from nucleotide 183 to nucleotide 884;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb28_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb28_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb28_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb28_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:32;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:31.

41. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:32;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vb28_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

42. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:33;
- (b) the nucleotide sequence of SEQ ID NO:33 from nucleotide 42 to nucleotide 206;
- (c) the nucleotide sequence of SEQ ID NO:33 from nucleotide 111 to nucleotide 206;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb29_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb29_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb29_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb29_1 deposited with the ATCC under accession number PTA-362;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:34;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:33.

43. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:34;

(b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb29_1 deposited with the ATCC under accession number PTA-362; the protein being substantially free from other mammalian proteins.

44. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 17 to nucleotide 253;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 98 to nucleotide 253;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb30_1 deposited with the ATCC under accession number PTA-362;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vb30_1 deposited with the ATCC under accession number PTA-362;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb30_1 deposited with the ATCC under accession number PTA-362;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb30_1 deposited with the ATCC under accession number PTA-362;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:36;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:35.

45. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:36;

(b) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb30_1 deposited with the ATCC under accession number PTA-362; the protein being substantially free from other mammalian proteins.

46. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:37;
- (b) the nucleotide sequence of SEQ ID NO:37 from nucleotide 68 to nucleotide 424;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vc67_1 deposited with the ATCC under accession number PTA-362;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc67_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:38;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:37.

47. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:38;
- (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc67_1 deposited with the ATCC under accession number PTA-362; the protein being substantially free from other mammalian proteins.

48. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:39;
- (b) the nucleotide sequence of SEQ ID NO:39 from nucleotide 103 to nucleotide 261;
- (c) the nucleotide sequence of SEQ ID NO:39 from nucleotide 154 to nucleotide 261;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vf4_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vf4_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vf4_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vf4_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:40;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:39.

49. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:40;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vf4_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

50. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:41;
- (b) the nucleotide sequence of SEQ ID NO:41 from nucleotide 1575 to nucleotide 3038;
- (c) the nucleotide sequence of SEQ ID NO:41 from nucleotide 1650 to nucleotide 3038;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vg3_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vg3_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vg3_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vg3_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:42;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:41.

51. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:42;

(b) a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42; and

(c) the amino acid sequence encoded by the cDNA insert of clone vg3_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

52. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:43;

(b) the nucleotide sequence of SEQ ID NO:43 from nucleotide 2112 to nucleotide 2363;

(c) the nucleotide sequence of the full-length protein coding sequence of clone vo2_1 deposited with the ATCC under accession number PTA-362;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo2_1 deposited with the ATCC under accession number PTA-362;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:44;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:43.

53. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:44;

(b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo2_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

54. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:45;

(b) the nucleotide sequence of SEQ ID NO:45 from nucleotide 36 to nucleotide 707;

(c) the nucleotide sequence of SEQ ID NO:45 from nucleotide 393 to nucleotide 707;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo3_1 deposited with the ATCC under accession number PTA-362;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo3_1 deposited with the ATCC under accession number PTA-362;

- (f) the nucleotide sequence of a mature protein coding sequence of clone vo3_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo3_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:46;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:45.

55. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo3_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

56. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:47;
- (b) the nucleotide sequence of SEQ ID NO:47 from nucleotide 74 to nucleotide 295;

- (c) the nucleotide sequence of SEQ ID NO:47 from nucleotide 134 to nucleotide 295;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo5_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo5_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo5_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo5_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:48;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:47.

57. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:48;
- (b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vo5_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

58. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:49;
- (b) the nucleotide sequence of SEQ ID NO:49 from nucleotide 45 to nucleotide 383;
- (c) the nucleotide sequence of SEQ ID NO:49 from nucleotide 312 to nucleotide 383;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo6_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo6_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo6_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo6_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:50;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:49.

59. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
- (b) a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vo6_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

60. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:51;
- (b) the nucleotide sequence of SEQ ID NO:51 from nucleotide 186 to nucleotide 1739;
- (c) the nucleotide sequence of SEQ ID NO:51 from nucleotide 288 to nucleotide 1739;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo9_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo9_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo9_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo9_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:51.

61. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:52;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo9_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

62. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:53;
- (b) the nucleotide sequence of SEQ ID NO:53 from nucleotide 440 to nucleotide 835;
- (c) the nucleotide sequence of SEQ ID NO:53 from nucleotide 632 to nucleotide 835;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo11_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo11_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo11_1 deposited with the ATCC under accession number PTA-366;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo11_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:54;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:53.

63. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:54;

(b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo11_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

64. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:55;

(b) the nucleotide sequence of SEQ ID NO:55 from nucleotide 72 to nucleotide 329;

(c) the nucleotide sequence of SEQ ID NO:55 from nucleotide 120 to nucleotide 329;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo12_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo12_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo12_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo12_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:56;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:55.

65. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:56;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo12_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

66. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:57;
- (b) the nucleotide sequence of SEQ ID NO:57 from nucleotide 227 to nucleotide 439;
- (c) the nucleotide sequence of SEQ ID NO:57 from nucleotide 287 to nucleotide 439;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo13_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo13_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo13_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo13_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:58;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:57.

67. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:58;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo13_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

68. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:59;
- (b) the nucleotide sequence of SEQ ID NO:59 from nucleotide 96 to nucleotide 341;
- (c) the nucleotide sequence of SEQ ID NO:59 from nucleotide 174 to nucleotide 341;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo14_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo14_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo14_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo14_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:60;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:59.

69. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:60;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo14_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

70. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:61;
- (b) the nucleotide sequence of SEQ ID NO:61 from nucleotide 90 to nucleotide 599;
- (c) the nucleotide sequence of SEQ ID NO:61 from nucleotide 165 to nucleotide 599;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo15_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo15_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo15_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo15_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:62;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:61.

71. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:62;

(b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo15_1 deposited with the ATCC under accession number PTA-366;
the protein being substantially free from other mammalian proteins.

72. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:63;

(b) the nucleotide sequence of SEQ ID NO:63 from nucleotide 209 to nucleotide 451;

(c) the nucleotide sequence of SEQ ID NO:63 from nucleotide 398 to nucleotide 451;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo16_1 deposited with the ATCC under accession number PTA-366;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo16_1 deposited with the ATCC under accession number PTA-366;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo16_1 deposited with the ATCC under accession number PTA-366;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo16_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:64;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:63.

73. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:64;

(b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo16_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

74. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:65;
- (b) the nucleotide sequence of SEQ ID NO:65 from nucleotide 31 to nucleotide 231;
- (c) the nucleotide sequence of SEQ ID NO:65 from nucleotide 97 to nucleotide 231;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo18_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo18_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo18_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo18_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:65.

75. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:66;

- (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo18_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

76. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:67;
- (b) the nucleotide sequence of SEQ ID NO:67 from nucleotide 23 to nucleotide 736;
- (c) the nucleotide sequence of SEQ ID NO:67 from nucleotide 83 to nucleotide 736;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo19_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo19_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo19_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo19_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:68;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:67.

77. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:68;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo19_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

78. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:69;
- (b) the nucleotide sequence of SEQ ID NO:69 from nucleotide 104 to nucleotide 1399;
- (c) the nucleotide sequence of SEQ ID NO:69 from nucleotide 158 to nucleotide 1399;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo22_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo22_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo22_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo22_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:70;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:69.

79. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:70;

(b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo22_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

80. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:71;

(b) the nucleotide sequence of SEQ ID NO:71 from nucleotide 174 to nucleotide 1595;

(c) the nucleotide sequence of the full-length protein coding sequence of clone vo23_1 deposited with the ATCC under accession number PTA-366;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo23_1 deposited with the ATCC under accession number PTA-366;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:72;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:71.

81. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:72;

(b) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo23_1 deposited with the ATCC under accession number PTA-366;
the protein being substantially free from other mammalian proteins.

82. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:73;

(b) the nucleotide sequence of SEQ ID NO:73 from nucleotide 129 to nucleotide 311;

(c) the nucleotide sequence of SEQ ID NO:73 from nucleotide 195 to nucleotide 311;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo24_1 deposited with the ATCC under accession number PTA-366;

- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo24_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo24_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo24_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:74;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:73.

83. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo24_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

84. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:75;
- (b) the nucleotide sequence of SEQ ID NO:75 from nucleotide 73 to nucleotide 798;
- (c) the nucleotide sequence of SEQ ID NO:75 from nucleotide 142 to nucleotide 798;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo25_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo25_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo25_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo25_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:75.

85. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;

- (b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo25_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

86. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:27;
- (b) the nucleotide sequence of SEQ ID NO:27 from nucleotide 26 to nucleotide 307;
- (c) the nucleotide sequence of SEQ ID NO:27 from nucleotide 101 to nucleotide 307;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo26_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo26_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo26_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo26_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:27.

87. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo26_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

88. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:79;
- (b) the nucleotide sequence of SEQ ID NO:79 from nucleotide 43 to nucleotide 228;
- (c) the nucleotide sequence of SEQ ID NO:79 from nucleotide 94 to nucleotide 228;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp23_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp23_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp23_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp23_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:80;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:79.

89. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:80;

(b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp23_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

90. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:81;

(b) the nucleotide sequence of SEQ ID NO:81 from nucleotide 245 to nucleotide 427;

(c) the nucleotide sequence of SEQ ID NO:81 from nucleotide 308 to nucleotide 427;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq7_1 deposited with the ATCC under accession number PTA-368;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq7_1 deposited with the ATCC under accession number PTA-368;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq7_1 deposited with the ATCC under accession number PTA-368;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq7_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:82;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:81.

91. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:82;

(b) a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq7_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

92. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:83;
- (b) the nucleotide sequence of SEQ ID NO:83 from nucleotide 119 to nucleotide 475;
- (c) the nucleotide sequence of SEQ ID NO:83 from nucleotide 185 to nucleotide 475;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq8_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq8_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq8_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq8_1 deposited with the ATCC under accession number PTA-368;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:84;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:83.

93. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:84;

- (b) a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vq8_1 deposited with the ATCC under accession number PTA-368; the protein being substantially free from other mammalian proteins.

94. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:85;
- (b) the nucleotide sequence of SEQ ID NO:85 from nucleotide 90 to nucleotide 323;
- (c) the nucleotide sequence of SEQ ID NO:85 from nucleotide 141 to nucleotide 323;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq9_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq9_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq9_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq9_1 deposited with the ATCC under accession number PTA-368;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:86;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:85.

95. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:86;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vq9_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

96. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:87;
- (b) the nucleotide sequence of SEQ ID NO:87 from nucleotide 18 to nucleotide 452;
- (c) the nucleotide sequence of SEQ ID NO:87 from nucleotide 72 to nucleotide 452;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq10_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq10_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq10_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq10_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:88;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:87.

97. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:88;

(b) a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq10_1 deposited with the ATCC under accession number PTA-368; the protein being substantially free from other mammalian proteins.

98. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:89;

(b) the nucleotide sequence of SEQ ID NO:89 from nucleotide 196 to nucleotide 378;

(c) the nucleotide sequence of SEQ ID NO:89 from nucleotide 262 to nucleotide 378;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq13_1 deposited with the ATCC under accession number PTA-368;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq13_1 deposited with the ATCC under accession number PTA-368;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq13_1 deposited with the ATCC under accession number PTA-368;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq13_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:90;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:89.

99. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:90;

(b) a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq13_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

100. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:91;
- (b) the nucleotide sequence of SEQ ID NO:91 from nucleotide 35 to nucleotide 718;
- (c) the nucleotide sequence of SEQ ID NO:91 from nucleotide 173 to nucleotide 718;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq16_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq16_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq16_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq16_1 deposited with the ATCC under accession number PTA-368;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:92;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:91.

101. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:92;

- (b) a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vq16_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

102. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:93;
- (b) the nucleotide sequence of SEQ ID NO:93 from nucleotide 1 to nucleotide 762;
- (c) the nucleotide sequence of SEQ ID NO:93 from nucleotide 70 to nucleotide 762;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq19_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq19_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq19_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq19_1 deposited with the ATCC under accession number PTA-368;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:94;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:93.

103. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:94;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vq19_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

104. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:95;
- (b) the nucleotide sequence of SEQ ID NO:95 from nucleotide 106 to nucleotide 792;
- (c) the nucleotide sequence of SEQ ID NO:95 from nucleotide 172 to nucleotide 792;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq20_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq20_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq20_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq20_1 deposited with the ATCC under accession number PTA-368;

- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:96;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:95.

105. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:96;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vq20_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

106. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:97;
- (b) the nucleotide sequence of SEQ ID NO:97 from nucleotide 40 to nucleotide 315;
- (c) the nucleotide sequence of SEQ ID NO:97 from nucleotide 124 to nucleotide 315;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq21_1 deposited with the ATCC under accession number PTA-368;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq21_1 deposited with the ATCC under accession number PTA-368;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq21_1 deposited with the ATCC under accession number PTA-368;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq21_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:98;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:97.

107. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:98;

(b) a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq21_1 deposited with the ATCC under accession number PTA-368;
the protein being substantially free from other mammalian proteins.

108. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:99;
- (b) the nucleotide sequence of SEQ ID NO:99 from nucleotide 70 to nucleotide 699;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vr2_1 deposited with the ATCC under accession number PTA-368;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vr2_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:100;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:99.

109. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:100;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vr2_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

110. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:101;
- (b) the nucleotide sequence of SEQ ID NO:101 from nucleotide 170 to nucleotide 394;
- (c) the nucleotide sequence of SEQ ID NO:101 from nucleotide 227 to nucleotide 394;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone PTA-1075 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc69_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc69_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc69_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:101.

111. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:102;

- (b) a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vc69_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins.

112. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:103;
- (b) the nucleotide sequence of SEQ ID NO:103 from nucleotide 43 to nucleotide 198;
- (c) the nucleotide sequence of SEQ ID NO:103 from nucleotide 85 to nucleotide 198;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc71_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc71_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc71_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc71_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:104;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:103.

113. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vc71_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins.

114. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:105;
- (b) the nucleotide sequence of SEQ ID NO:105 from nucleotide 260 to nucleotide 1552;
- (c) the nucleotide sequence of SEQ ID NO:105 from nucleotide 335 to nucleotide 1552;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo27_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo27_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo27_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo27_1 deposited with the ATCC under accession number PTA-1075;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:106;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:105.

115. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:106;

(b) a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo27_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins.

116. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:107;

(b) the nucleotide sequence of SEQ ID NO:107 from nucleotide 15 to nucleotide 320;

(c) the nucleotide sequence of SEQ ID NO:107 from nucleotide 72 to nucleotide 320;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo31_1 deposited with the ATCC under accession number PTA-1075;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo31_1 deposited with the ATCC under accession number PTA-1075;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo31_1 deposited with the ATCC under accession number PTA-1075;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo31_1 deposited with the ATCC under accession number PTA-1075;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:108;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:107.

117. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:108;

(b) a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo31_1 deposited with the ATCC under accession number PTA-1075; the protein being substantially free from other mammalian proteins.

118. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:109;
- (b) the nucleotide sequence of SEQ ID NO:109 from nucleotide 38 to nucleotide 1255;
- (c) the nucleotide sequence of SEQ ID NO:109 from nucleotide 86 to nucleotide 1255;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo32_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo32_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo32_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo32_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:110;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:109.

119. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:110;

- (b) a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo32_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins.

120. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:111;
- (b) the nucleotide sequence of SEQ ID NO:111 from nucleotide 80 to nucleotide 1276;
- (c) the nucleotide sequence of SEQ ID NO:111 from nucleotide 131 to nucleotide 1276;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo33_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo33_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo33_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo33_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:112;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:111.

121. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:112;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone vo33_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins.

122. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:113;
- (b) the nucleotide sequence of SEQ ID NO:113 from nucleotide 202 to nucleotide 429;
- (c) the nucleotide sequence of SEQ ID NO:113 from nucleotide 292 to nucleotide 429;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq23_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq23_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq23_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq23_1 deposited with the ATCC under accession number PTA-1075;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:114;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:113.

123. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:114;

(b) a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq23_1 deposited with the ATCC under accession number PTA-1075; the protein being substantially free from other mammalian proteins.

124. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:115;

(b) the nucleotide sequence of SEQ ID NO:115 from nucleotide 37 to nucleotide 1113;

(c) the nucleotide sequence of SEQ ID NO:115 from nucleotide 88 to nucleotide 1113;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq24_1 deposited with the ATCC under accession number PTA-1075;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq24_1 deposited with the ATCC under accession number PTA-1075;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq24_1 deposited with the ATCC under accession number PTA-1075;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq24_1 deposited with the ATCC under accession number PTA-1075;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:116;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:115.

125. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:116;

(b) a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq24_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins.

126. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:117;
- (b) the nucleotide sequence of SEQ ID NO:117 from nucleotide 40 to nucleotide 207;
- (c) the nucleotide sequence of SEQ ID NO:117 from nucleotide 103 to nucleotide 207;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq26_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq26_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq26_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq26_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:118;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:117.

127. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:118;

- (b) a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vq26_1 deposited with the ATCC under accession number PTA-1075; the protein being substantially free from other mammalian proteins.